Naval Research Laboratory

Washington, DC 20375-5000



NRL Memorandum Report 6268

AD-A200 187

The Electron — Water Vapor (H₂O) Collision Cross Sections

A. W. ALI

Plasma Physics Division



August 26, 1988

Approved for public release; distribution unlimited.

| SECURITY CLASSIFICATION OF THIS PAGE | | | | | |
|---|--------------------------------------|-----------------------------|---------------------------------|---|--|
| REPORT | DOCUMENTATIO | N PAGE | | Form Approved OMB No 0704 0188 | |
| 1a REPORT SECURITY CLASSIFICATION UNCLASSIFIED | _ | 16 RESTRICTIVE MARKINGS | | | |
| Za SECURITY CLASSIFICATION AUTHORITY | 2a SECURITY CLASSIFICATION AUTHORITY | | V AVAILABILITY OF | REPOPT | |
| 26 DECLASSIFICATION DOWNGRADING SCHED | JL E | Approved f unlimited. | | lease; distribution | |
| 4 PERFORMING ORGANIZATION REPORT NUMB | ER(S) | 5 MONITORING | ORGANIZATION RE | PORT NUMBER(S) | |
| NRL Memorandum Report 6268 | | ľ | | | |
| 6a NAME OF PERFORMING ORGANIZATION | 7a NAME OF M | ONITORING OPGAN | 7A1 ON | | |
| Naval Research Laboratory 6c ADDRESS (City State, and ZIP Code) | 7h ADDRESS (C. | ty State and ZIP Co | ado) | | |
| Washington, DC 20375-5000 | | 78 2001111111 | ly state and zir co | | |
| 8a NAME OF FUNDING SPONSORING ORGANIZATION | 8b OFFICE SYMBOL (If applicable) | 9 PROCUREMEN | T INSTRUMENT DE | NTFICATION N. MBER | |
| DARPA | <u> </u> | 10. 50. 865.05 | 5. 1.0.1.6.1.40.506 | | |
| 8c ADDRESS (City State and ZIP Code) | | PROGRAM | FUNDING NUMBERS | TASK WORK JAIT | |
| Arlington, VA 22209 | | 62707E | NO N60921- 87-WX-0072 | ACCESSION NO 4395, A63 DN680-415 | |
| 11 TITLE (Include Security Classification) | | - L | -t | | |
| The Electron-Waters Vapor (H | O) Collision Cr | coss Section | S | | |
| 12 PERSONAL AUTHOR(S) Ali, A.W. | | | | | |
| 13a TYPE ÖF REPORT 13b TIME C Interim FROM | OVERED 10 | 14 DATE OF REPO 1988 Aug | ORT (Year, Month, D ust 26 | 15 PAGE COUNT | |
| 16 SUPPLEMENTARY NOTATION | | | | | |
| 17 COSATI CODES | 18 SUBJECT TEPMS (| Continue on revers | se if necessary and | identify by block number) | |
| FIELD GROUF SUB-GROUP | 1 H ₂ O | Commisc 311 12 (2) | Electron i | mpact cross sections | |
| | lonoization ≱Inclastic coll | isione . | → Elastic co Momentum t | × | |
| 19 ABSTPAC* (Continue on reverse if necessary | | | Monie recting t | ransfert A. | |
| A set for electron-H ₂ O collisi The set comprises the total, ionization rotational, elastic, and momentum tra | on, dissociative ioni | zation, dissocia | er a wide rangative excitation. | e of electron energies. electronic, vibrational, | |
| 20 DISTRIBUTION AVAILABILITY OF ABSTRACT XX UNCLASSIFIED UNLIMITED | RPT DTIC USERS | 21 ABSTRACT SE UNCLASSIF | CURITY CLASSICAT LED | 7-(30, | |
| 22a NAME OF RESPONSIBLE INDIVIDUAL | CONTROL ON FRA | | Include Area Code) | 22c 000G S7780. Code 4700,1 | |
| _ A.W. Ali | (404) 101- | <u> </u> | Conc arount | | |

Previous editions are obsolete

DD Form 1473, JUN 86

SECURITY CLASSIFICATION OF THIS PAUL

CONTENTS

| 1. | INTRODUCTION | 1 |
|------|---|----------------|
| 2. | THE IONIZATION OF H ₂ 0 | 2 |
| 3. | THE DISSOCIATION OF H ₂ 0 | 3 |
| 4. | THE EXCITATION OF H ₂ 0 | 6 |
| | | 6 10 10 |
| 5. | THE TOTAL, ELASTIC AND MOMENTUM TRANSFER CROSS SECTIONS | 1 3 |
| | 5.2 The Elastic Cross Section | 13 13 17 |
| 6. | THE DISSOCIATIVE ATTACHMENT CROSS SECTION | 17 |
| REFE | RENCES | 21 |
| DIST | RIBUTION | 31 |





THE ELECTRON—WATER VAPOR (H₂O) COLLISION CROSS SECTIONS

1. INTRODUCTION

Water vapor (H_20) is one of the minor constituents of the atmosphere and it plays an important role in the deionization processes of the partially ionized air. It controls the air conductivity, especially that of the unheated air. Therefore, the role of H_20 in modifying the electron velocity distribution in wet air is required for characterizations of partially ionized wet air. This requires the solution of a Boltzmann equation with an appropriate set of electron-molecule scattering cross sections.

This report, accordingly, presents a set of up to date cross sections for the electron - H_2O collisions over a wide range of electron energies.

Manuscript approved May 12, 1988.

2.0 THE IONIZATION OF H₂0

The electron impact ionization of ${\rm H_20}$ results in the ionization of ${\rm H_2^20}$ as well as in several dissociative ionization channels. These are:

$$e + H_2O \rightarrow H_2O^+ + 2e$$
 (1)

$$e + H_2O \rightarrow OH^+ + H + 2e$$
 (2)

$$e + H_2O \rightarrow 0^+ + H_2 + 2e$$
 (3)

$$e + H_2O \rightarrow H_2^+ + O + 2e$$
 (4)

$$e + H_2O \rightarrow H^+ + OH + 2e$$
 (5)

One of the very early measurements by Mann, et al¹, in 1940, of the $\rm H_2O$ ionization cross sections, gave only the relative ion abundancies generated by electrons with energy of 100 eV. These values were 100, 23.2, 2.0, 0.07 and 5.0 for the relative productions of $\rm H_2O^+$, $\rm OH^+$, $\rm O^+$, and $\rm H^+_2$ and $\rm H^+$, respectively. However, the measurements of Schutten, et al², some 26 years later provided partial ionization cross sections for the above processes for electron energies from 20 to 1000 eV. It also provided the relative abundancies of the ions produced by the impact of electrons with energy of 100 eV.

These are 100, 25.8, 2.65, 0.3 and 29.6 for H₂0⁺, 0H⁺, 0⁺, H₂⁺ and H⁺, respectively. These results differ sharply from those of Mann et al¹, especially for the productions of H₂⁺ and H⁺. The total and partial ionization cross sections of H₂0 are shown in Fig. 1 and are also given in Table 1, based on data from Ref. 2, where it is obvious that the dominant contributions to the total ionization are from H₂0⁺, 0H⁺ and H⁺. These values are from threshold energies to 1000 eV; however, for higher impact energies, the ionization cross section can be obtained by using the following equation³.

$$\sigma_{i} = A_{i} \frac{4\pi a_{0}^{2} R^{2}}{I_{i}E'} \left[log \frac{4C_{i}E'}{I_{i}} (1-\beta^{2}) - \beta^{2} \right]$$
 (6)

where $E' = \frac{1}{2} \text{ mc}^2 \beta^2$, A_i and C_i are adjustable parameters, a_o is the Bohr radius and R is the Rydbeg constant (13.6 eV).

3. THE DISSOCIATION OF H,0

The dissociation of H₂0 has many channels. These are:

$$e + H_2O \rightarrow OH + H + e$$
 (7)

$$e + H_2 0 \rightarrow H_2 + 0 + e$$
 (8)

$$e + H_2 0 \rightarrow H + H + 0 + e$$
 (9)

No absolute measurements of these cross sections exist. However, absolute emission cross sections for the dissociative excitation

processes have been measured. These dissociative excitations are; the OH ($A^2\Sigma$) excitation which emits radiation in the 2810 – 3500Å, the La, Ha, Hb, Hy and Hb emissions from the hydrogen atom and the 1302Å, 7774 Å and 8447Å emissions from the oxygen atom. The emission cross sections of OH ($A^2\Sigma$), Hb, O(8447Å) and O(7774Å) have been measured by Beenakker, et al4, in the energy range of 20 – 1000 eV. Furthermore, the emission cross sections for Ha, Hy, and Ha have been measured at 300 eV and are normalized to that of Hb. The measurements of these emissions at 100, 200, 500 and 800 eV indicate4 the same energy dependence, as that of Hb.

TABLE I
TOTAL AND PARTIAL IONIZATION CROSS SECTIONS

ENERGY (eV)

| | H ₂ 0+ | 0H+ | 0+ | 0++ | H ₂ + | H+ | TOTAL |
|------|-------------------|----------|----------|----------|------------------|----------|-----------|
| 20 | 4.2(-17)* | 3.0(-18) | 9.0(-20) | | 1.8(-19) | 7.0(-19) | 4.6(-17) |
| 30 | 7.6(-17) | 1.3(-17) | 3.7(-19) | | 2.2(-19) | 7.8(-18) | 9.7(17) |
| 50 | 1.07(-16) | 2.4(-17) | 1.7(-18) | | 3.0(-19) | 2.4(-17) | 1.57(-16) |
| 100 | 1.32(-16) | 3.4(-17) | 3.5(-18) | | 3.9(-19) | 3.9(-17) | 2.10(-16) |
| 150 | 1.16(-16) | 3.1(-17) | 3.6(-18) | 6.0(-20) | 3.6(-19) | 3.7(-17) | 1.89(-16) |
| 200 | 1.08(-16) | 2.8(-17) | 3.5(-18) | 1.4(-19) | 2.8(-19) | 3.6(-17) | 1.75(-16) |
| 500 | 6.4(-17) | 1.7(-17) | 1.9(-18) | 1.3(-19) | 1.8(-19) | 2.3(-17) | 1.06(-16) |
| 700 | 5.0(-17) | 1.4(-17) | 1.4(-18) | 9.0(-20) | 1.6(-19) | 1.7(-17) | 8.3(-17) |
| 1000 | 4.1(-17) | 1.1(-17) | 1.1(-18) | 6.0(-20) | 8.0(-20) | 1.3(-17) | 6.5(-17) |
| 1500 | 2.9(-17) | 8.0(-18) | 8.0(-19) | | | 1.1(-17) | 4.9(-17) |
| 2000 | 2.3(-17) | 6.0(-18) | 6.0(-19) | | | 8.0(-18) | 3.7(-17) |

 $[\]star 4.2(-17)$ implies $4.2 \times 10^{-17} \text{cm}^2$

These cross sections, based on the measurements of Beenakker, et al⁴, are given in Table II and are shown in Figure 2. The excitation of the 1304Å line of oxygen from the dissocation product has been measured by Lawrence⁵ and the absolute emission cross section of Ly has been measured by Vroom and de Heer⁶. The data for the last two cross sections is given in Table III and are also shown in Fig. 2.

4. THE EXCITATION OF H₂0

4.1 The Electronic Excitations

There exist no measured cross sections for the electronic excitations for $\mathrm{H}_2\mathrm{O}$. However, high energy electron scattering spectra have been obtained by Lassettre and co-worker⁷⁻¹⁰. The generalized oscillator strengths obtained by these workers can be utilized¹¹ to obtain the excitation cross sections through the appropriate relations, especially for the optically allowed transitions. Trajmar, et al,^{12,13}, have detected two excitations in $\mathrm{H}_2\mathrm{O}$ at 4.5 and 9.8 eV with features appropriate to triplet - singlet transitions. However, to date, theoretical calculations (See Ref. 13, 14) indicate no electronic excitations for states below 6 eV. Yousfi, et al¹⁴, give the cross sections for two $\mathrm{H}_2\mathrm{O}$ excitations with thresholds at 7.5 and 11.8 eV. These cross sections are shown in Figure 3 and are also given in Table IV.

| Energy (eV) | OH (A ² Σ) | Нα | нв | Η σ | н 8 | 0 8447Å | 0 777 4 Å |
|----------------|-------------------------|-------|------|-------|-------|------------|---------------------|
| 40 | 66.3 | 15.82 | 3.06 | 1.132 | 0.428 | 1.40 | 0.95 |
| 50 | 60.3 | 20.88 | 4.04 | 1.49 | 0.56 | 1.88 | 1.16 |
| 60 | 56.6 | 27.14 | 5.25 | 1.94 | 0.73 | 2.30 | 1.29 |
| 70 | 53.5 | 30.45 | 5.89 | 2.18 | 0.82 | 2.54 | 1.33 |
| 80 | 50.8 | 32.31 | 6.25 | 2.31 | 0.87 | 2.70 | 1.32 |
| 90 | 48.3 | 33.09 | 6.4 | 2.37 | 0.89 | 2.8 | 1.28 |
| 100 | 46.4 | 33.14 | 6.41 | 2.37 | 0.89 | 2.86 | 1.26 |
| 120 | 43.2 | 31.69 | 6.13 | 2.27 | 0.86 | 2.87 | 1.17 |
| 140 | 40.0 | 30.14 | 5.83 | 2.16 | 0.82 | 2.59 | 1.02 |
| 170 | 36.4 | 26.98 | 5.22 | 1.93 | 0.73 | 2.53 | 0.80 |
| 200 | 33.9 | 25.13 | 4.86 | 1.79 | 0.68 | 2.32 | 0.73 |
| 250 | 29.7 | 21.61 | 4.18 | 1.55 | 0.58 | 1.9 | 0.54 |
| 300 | 26.6 | 19.02 | 3.68 | 1.36 | 0.51 | 1.67 | 0.46 |
| 400 | 22.4 | 15.09 | 2.92 | 1.08 | 0.41 | 1.30 | 0.36 |
| 450 | 20.6 | 13.54 | 2.62 | 0.97 | 0.36 | 1.13 | 0.35 |
| 500 | 19.6 | 12.35 | 2.39 | 0.88 | 0.33 | 1.05 | 0.28 |
| 600 | 17.9 | 10.59 | 2.05 | 0.76 | 0.28 | 0.89 | 0.24 |
| 700 | 15.8 | 9.41 | 1.82 | 0.67 | 0.25 | 0.78 | 0.17 |
| 800 | 14.9 | 8.32 | 1.61 | 0.59 | 0.22 | 0.68 | 0.17 |
| 900 | 13.1 | 7.6 | 1.47 | 0.54 | 0.20 | 0.58 | 0.16 |
| 1000 | 12.1 | 7.13 | 1.38 | 0.51 | 0.19 | 0.53 | 0.14 |

TABLE III
EMISSION CROSS SECTIONS

| Energy (eV) | Lγ | 1304Å |
|-------------|-----------|----------|
| 40 | | 1.3(-19) |
| 50 | 2.54(-17) | 2.8(-19) |
| 60 | 2.60(-17) | 3.3(-19) |
| 80 | 2.61(-17) | 3.7(-19) |
| 100 | 2.58(-17) | 3.7(-19) |
| 150 | 2.14(-17) | 2.7(-19) |
| 200 | 1.87(-17) | 2.3(-19) |

TABLE IV ${\tt EXCITATION} \ \, {\tt CROSS} \ \, {\tt SECTIONS} \ \, {\tt OF} \ \, {\tt H}_2{\tt O}$

| E(ev) | Eth=7.1 eV | Eth=11.8 eV |
|-------|------------|-------------|
| 7.1 | 0 | |
| 11.8 | 0 | |
| 15 | 0.12(-16) | |
| 16 | | 0.078(-16) |
| 20 | 0.2(-16) | 0.135(-16) |
| 30 | 0.27(-16) | 0.180(-16) |
| 40 | 0.32(-16) | 0.213(-16) |
| 55 | 0.33(-16) | 0.225(-16) |
| 80 | 0.30(-16) | 0.202(-16) |
| 100 | 0.25(-16) | 0.168(~16) |
| 120 | 0.20(-16) | 0.135(-16) |
| 2.40 | 0.15(-16) | 0.101(-16) |
| 180 | 0.12(-16) | 0.078(-16) |
| 260 | 0.07(-16) | 0.045(-16) |
| 320 | 0.033(-16) | 0.022(-16) |
| 400 | 0.017(-16) | 0.011(-16) |

4.2 The Vibrational Cross Sections

The vibrational excitation of H_2O consists of the stretching modes ν_1 , ν_3 , the bending mode ν_2 , their harmonics and sums of these frequencies (see e.g. Ref. 10). However, the emission measurements of Ref. 10 and the most recent cross section measurements¹⁵, ¹⁶, indicate that ν_1 , ν_2 , and ν_3 are the strongest vibrational transitions in H_2O . Furthermore, ν_1 and ν_3 are blended together and are not separated experimentally. The cross section for ν_2 and (ν_1, ν_3) based on measurements of Rhor¹⁵ and Seng and Linder¹⁶ are shown in Fig. 4 and are also given in Table V.

4.3 The Rotational Cross Sections

Experimental data on the rotational excitation of $\mathrm{H}_2\mathrm{O}$ is limited¹⁷ to the differential scattering cross sections at impact energies of 2.4 and 6.0 eV. These measurements by Jung, et al¹⁷, are for the excitation and de-excitation processes $(\Delta j = \pm 1)$. Jung, et al¹⁷, have also measured the differential cross sections for the elastic process at the two energies mentioned earlier.

These differential cross sections have been integrated by Danjo and Nishimura¹⁸ to provide the total elastic scattering cross section at 2.14 and 6.0 eV. These values are in good agreement with the total elastic scattering cross section measured by Danjo and Nishimura¹⁸. This provides reasonable confidence in the magnitude of the differential cross sections for rotational excitation . Furthermore, Jain and Thompson¹⁹ have calculated the rotational excitations of $\rm H_2O$ in the range of 1 - 10 eV. The calculated differential cross sections, however, are at least a factor of 3 larger in comparison with the measured values of Jung, et al¹⁷, at 2.14 and 6.0 eV. These results are shown in Figure 5 along with the proposed rotational cross section by Yousfi, et al¹⁴, based on swarm data and solution of the Boltzmann equation.

| Energy (eV) | (v_1, v_3) | (v ₂) |
|-------------|--------------|-------------------|
| 0.2 | 0 | 0 |
| 0.3 | 2.0(-19) | 3.0(-17) |
| 0.4 | 3.0(-18) | 9.5(-17) |
| 0.5 | 3.0(-17) | 6.0(-17) |
| 0.6 | 1.4(-16) | 4.5(-17) |
| 0.7 | 2.0(-16) | 3.0(-17) |
| 0.8 | 1.35(-16) | 2.5(-17) |
| 0.9 | 9.5(-17) | 1.7(-17) |
| 1.0 | 7.0(-17) | 1.35(-17) |
| 1.5 | 1.95(-17) | 7.5(-18) |
| 1.6 | 1.75(-17) | 7.55-18) |
| 1.8 | 1.75(-17) | 8.0(-18) |
| 2.0 | 1.95(-17) | 8.5(-18) |
| 2.2 | 2.05(-17) | 9.0(-18) |
| 2.4 | 2.15(-17) | 9.5(-18) |
| 2.6 | 2.20(-17) | 1.0(-17) |
| 2.8 | 2.30(-17) | 1.05(-17) |
| 3.0 | 2.35(-17) | 1.1(-17) |
| 3.5 | 2.45(-17) | 1.25(-17) |
| 4.0 | 2.60(-17) | 1.3(-17) |
| 5.0 | 2.80(-17) | 1.4(17) |
| 6.0 | 3.00(-17) | 1.5(-17) |
| 7.0 | 3.00(-17) | 1.5(-17) |
| 8.0 | 3.00(-17) | 1.4(-17) |
| 9.0 | 2.60(-17) | 1.2(-17) |
| 10 | 2.10(-17) | 9.5(-18) |

5. THE TOTAL, ELASTIC AND MOMENTUM TRANSFER CROSS SECTIONS

5.1 The Total Scattering Cross Section

The total scattering cross section of electrons in H₂0 has been measured from 0.5 to 1000 eV. The earliest measurement is that of Brüche²⁰ (1929) for impact energies of 1.5 to 40 eV. The other measurements are very recent and have been carried out from 1981 to 1987. These are: Sokolov²¹ (1981) in the energy range of 0.5 - 7 eV, Sueoka, et al²², (1986) from 1.0 to 80 eV, Szmytkowski, et al²³, and Szmytkowski²⁴ (1987) in the energy range of 0.5 - 1000 eV, and those of Seng and Linder (See Ref. 23) from 0.5 - 10 eV. These measurements are shown in Figure (6) and are given in Table (VI)

5.2 The Elastic Cross Section

The electron elastic scattering cross section in $\rm H_2O$ has been measured by Danjo, et al¹⁸, in the range of 4 to 200 eV and by Katase, et al²⁵, in the range of 100 - 1000 eV. These measured values are shown in Figure (7) along with the total scattering cross section of Szmytkowski²³. The numerical values of these elastic cross sections are given in Table (VIIA).

TABLE VI $\label{total} \mbox{TOTAL SCATTERING CROSS SECTION IN } (10^{-16}\mbox{cm}^2)$

| SZ | <u>B</u> | SL | so | su |
|------|--|--|--|--|
| | | | 3000 | |
| | | | 2000 | |
| | | | 1500 | |
| | | | 1100 | |
| | | | 800 | |
| | | | 600 | |
| 70.2 | | 81.1 | 125 | |
| 52.8 | | | 87 | |
| 40.2 | | 37.0 | 60 | 29.2 |
| 28.8 | 23.1 | | 40 | |
| 24.0 | 20.2 | 19.4 | 30 | 15.8 |
| 20.7 | 17.2 | 15.3 | 20 | |
| 19.5 | 16.1 | 13.9 | 16 | 12.2 |
| 19.8 | 16.3 | 14.2 | 13.3 | 11.9 |
| 20.0 | 16.5 | 15.3 | 11.8 | 12.5 |
| 20.7 | 16.8 | 16.7 | 10.8 | 12.6 |
| 20.9 | 17.4 | 17.7 | | 13.0 |
| 21.1 | 17.8 | 18.1 | | 13.2 |
| 20.9 | 18.1 | 18.1 | | 12.7 |
| 19.0 | 18.2 | | | 12.8 |
| | 70.2 52.8 40.2 28.8 24.0 20.7 19.5 19.8 20.0 20.7 20.9 21.1 | 70.2 52.8 40.2 28.8 23.1 24.0 20.2 20.7 17.2 19.5 16.1 19.8 16.3 20.0 16.5 20.7 16.8 20.9 17.4 21.1 17.8 20.9 18.1 | 70.2 81.1 52.8 37.0 28.8 23.1 24.0 20.2 19.4 20.7 17.2 15.3 19.5 16.1 13.9 19.8 16.3 14.2 20.0 16.5 15.3 20.7 16.8 16.7 20.9 17.4 17.7 21.1 17.8 18.1 20.9 18.1 18.1 | 3000 2000 1500 1100 800 600 70.2 81.1 52.8 87 40.2 37.0 60 28.8 23.1 40 24.0 20.2 19.4 30 20.7 17.2 15.3 20 19.5 16.1 13.9 16 19.8 16.3 14.2 13.3 20.0 16.5 15.3 11.8 20.7 16.8 16.7 10.8 20.9 17.4 17.7 21.1 17.8 18.1 20.9 18.1 18.1 |

TABLE VI TOTAL SCATTERING CROSS SECTION IN $(10^{-16} cm^2)$ (Continued)

| Energy (eV) | <u>SZ</u> | <u>B</u> | SL | <u>so</u> | <u>su</u> |
|-------------|-----------|-------------|----|-----------|-------------|
| 16 | 16.7 | 16.7 | | | 12.0 |
| 20 | 15.7 | 15.0 | | | 11.3 |
| 25 | 14.1 | 13.3 | | | 10.2 |
| 30 | 12.7 | 11.9 | | | 9.5 |
| 35 | 11.8 | 10.8 | | | 9.0 |
| 40 | 11.0 | 10 | | | 8.5 |
| 50 | 9.7 | | | | 7.5 |
| 60 | 8.2 | | | | 7.1 |
| 70 | 8.1 | | | | 6.7 |
| 80 | 7.5 | | | | 6.3 |
| 100 | 6.65 | | | | |
| 200 | 4.50 | | | | |
| 300 | 3.5 | | | | |
| 400 | 2.80 | | | | |
| 500 | 2.50 | | | | |
| 1000 | 1.43 | | | | |

Symbols: B = Ref 20

SL = See Ref 24

SO = Ref 21

SU = Ref 22 SZ = Refs 21 and 24

TABLE VIIA ELASTIC AND MOMENTUM TRANSFER CROSS SECTIONS

| Energy (eV) | <u>σ_e (D)</u> | $\underline{\sigma_{m}(D)}$ | <u>σ_e (K)</u> | $\sigma_{m}(K)$ |
|-------------|--------------------------|-----------------------------|----------------------------|-----------------|
| 4 | 1.04(-15) | 6.18(-16) | | |
| 5 | 1.05(-15) | 6.17(-16) | | |
| 6 | 1.06(-15) | 6.40(-16) | | |
| 8 | 1.11(-15) | 6.43(-16) | | |
| 10 | 1.2(-15) | 6.81(-16) | | |
| 15 | 9.93(-16) | 5.74(-16) | | |
| 20 | 8.21(-16) | 4.37(-16) | | |
| 30 | 5.72(-16) | 2.65(-16) | | |
| 40 | 4.68(-16) | 2.34(-16) | | |
| 50 | 3.97(-16) | 1.65(-16) | | |
| 60 | 3.34(-16) | 1.37(-16) | | |
| 80 | 2.63(-16) | 8.86(-17) | | |
| 100 | 2.37(-16) | 8.62(-17) | 2.98(-16) | 1.01(-16) |
| 150 | 1.78(-16) | 5.26(-17) | | |
| 200 | 1.50(-16) | 5.04(-17) | 2.11(-16) | 0.464(-16) |
| 300 | | | 1.56(-16) | 0.296(-16) |
| 400 | | | 1,32(-16) | 0.208(-16) |
| 500 | | | 1.04(-16) | 0.156(-16) |
| 700 | | | 0.819(-16) | 0.093(-16) |
| 1000 | | | 0.548(-16) | 0.0515(-16) |
| | | | | |

 $[\]sigma(D) = Ref 18$ $\sigma(K) = Ref 25$

5.3 The Momentum Transfer Cross Section

The electron momentum transfer cross section in H_2O has been measured by Danjo, et al¹⁸, in the energy range of 4.0 to 200 eV, and by Katase, et al²⁵, in the energy range of 100 to 1000 eV. These values are given in Table (VIIA). Furthermore, Pack, et al²⁶, have obtained the momentum transfer cross in H_2O for very low electron energies, i.e., for 0.003 to 0.08 eV through swarm data analysis. They give the following expression for the momentum transfer cross section, where E is

$$\sigma_{\rm m} = 10^{-14} \left[2.74E + 2.54(E)^{3/2} \right]^{-1}$$
 (10)

the electron energy in eV. Using Eq. (10) the numerical values of the cross section are given in Table VIIB along with data from Itikawa's compilation²⁷.

6. THE DISSOCIATIVE ATTACHMENT CROSS SECTION

Electron collisions with $\rm H_2O$ lead to the generation of negative ions in addition to all other processes discussed earlier. The negative ions are H and O and are produced by the process of the dissociative attachment.

$$e + H_2O \rightarrow \overline{H} + OH \tag{11}$$

$$e + H_2 0 \rightarrow \overline{0} + H_2 0$$
 (12)

TABLE VIIB

| ELECTRON ENERGY (eV) | MOMENTUM TRANS P | FER CROSS SECTION ELECTRON ENERGY (eV) | (cm²) <u>I</u> |
|-------------------------|------------------|--|-------------------|
| 0.01 | 3.34(-13) | 0.5 | 3.5(-15) |
| 0.02 | 1.61(-13) | 0.8 | 2.0(-15) |
| 0.04 | 7.69(-14) | 1.0 | 1.5(-15) |
| 0.06 | 4.96(-14) | 2.0 | 6.0(-15) |
| 0.08 | 3.61(-14) | .0 | 5.5(-15) |
| 0.1 | 2.82(-14) | 4.0 | 6.0(-15) |
| | | 5.0 | 6.8(-15) |
| | | 6.0 | 7.7(-15) |
| | | 7.0 | 7.9(-15) |
| | | 8.0 | 8.0(~15) |
| | | 10 | 7.5(-15) |

Symbols: P = Ref 26 I = Ref 27

Act of

Where the cross section for process (11) has been measured by Crompton and Christophorou²⁸ and Belic et al²⁹. The relative strength of process (12) is indicated28 to be one order of magnitude smaller than that of process (11). These cross sections are given in Table VIII. The onsets of process (11) and (12) occur at 5.7 and 7.5 eV and the peak of their cross sections are at 6.5 and 11.5 eV. respectively. Process (12), on the other hand, has two28 other peaks which are at 7 and 9 eV. These measurements have been made at low gas pressures, however, at higher pressures one observes OH formation which is due to ion-molecular reactions. One of the interesting observations of the H formation in process (11) is that OH is vibrationally excited²⁹. Belic, et al²⁸ have observed excited vibrational levels up to v = 7 and have resolved the excitations of individual levels as a function of the electron energy.

TABLE VIII
THE DISSOCIATIVE ATTACHMENT CROSS SECTION

| Electron Energy (eV) | H (Ref 28) | Electron Energy (eV) | H (Ref 27) | Electron Energy (eV) | 0 (Ref 27) |
|-------------------------|------------|-------------------------|------------|-------------------------|------------|
| 5.56 | 0.9(-18) | 5.6 | 0 | 5 | 0 |
| 5.87 | 2.0(-18) | 6.0 | 2.5(-18) | 6 | 3.9(~20) |
| 6.0 | 3.0(-18) | 6.5 | 6.9(-18) | 7 | 1.6(-19) |
| 6.25 | 5.0(-18) | 7.0 | 4.3(-18) | 8 | 3.9(-20) |
| 6.5 | 6.4(-18) | 7.5 | 1.6(-18) | 9 | 3.90(-19) |
| 6.75 | 5.2(-18) | 8.0 | 1.2(-18) | 10 | 2.7(-19) |
| 7.0 | 3.8(-18) | 8.5 | 1.25(-18) | 11 | 6.3(-19) |
| 7.25 | 2.5(-18) | 9.0 | 1.0(-18) | 11.5 | 6.9(-19) |
| 7.5 | 1.2(-18) | 9.5 | 0.5(-18) | 12 | 5.5(-19) |
| 7.7 | 0.8(-18) | 10.0 | 0.3(-18) | 13 | 1.8(-19) |
| | | 11 | 0.2(-18) | 14 | 7.0(-20) |
| | | | | 15 | 2.3(-20) |

REFERENCES

- M.M.Mann, A. Hustrulid and J. T. Tate, Phys. Rev. <u>58</u>, 340(1940)
- I. Schutten, F. J. De Heer, H. R. Moustafa, A. J. H. Boerboom and J. Kistemaker, J. Chem. Phys. 44, 3924 (1966)
- 3. R. D. Taylor and A. W. Ali, "Excitation and Ionization Cross Sections for Electron-Beam Energy Deposition in High Temperature Air", NRL Memorandum Report 6013 (1987)
- 4. C.I.M. Beenakker, F. J. De Heer, H. B. Krop and G. R. Mohlmann, Chem. Phys., 6, 445 (1974)
- 5. G. M. Lawrence, Phys. Rev. A2, 397 (1970)
- 6. D. A. Vroom and F. J. de Heer, J. Chem. Phys. 50, 1883 (1969)
- 7. E. N. Lassettre and E. R. White, J. Chem. Phys. <u>60</u>, 2460 (1974)
- 8. E. N. Lassettre and A. Skerbele, J. Chem. Phys. $\underline{60}$, 2464 (1974)
- 9. K. N. Klump and E. N. Lassettre, Can. J. Phys. <u>53</u>, 1825 (1975)
- 10. E. N. Lassettre, Can J. Chem. 47, 1733 (1969)
- J. J. Olivero, R. W. Stagat and A. E. S. Green, J. Geophys. Res. 77, 4797 (1972)
- 12. S. Trajmar, D. F. Register and A. Chutjean, Phys. Reports (Review Section of Physics Letters) 97, 219 (1983)
- 13. S. Trajmar, W. Williams and A. Kuppermann, J. Chem. Phys. $\underline{54}$, 2274 (1971)
- 14. M. Yousfi, N. Azzi, P. Segur, I. Gallimberti, and S. Stangherlin, "Electron-Molecule Collision Cross Sections and Electron Swarm Parameters in Some Atmosphere Gases", Centre de Physique Atomique de Toulouse, Private Communication 1987
- 15. K. Rohr, J. Phys. B 10, L735 (1977)
- G. Seng and F. Linder, J. Phys. B 9, 2539 (1976)

- K. Jung, Th. Antoni, R. Muller, K. H. Kochem and H. Ehrhardt, J. Phys. B <u>15</u>, 3535 (1982)
- A. Danjo and H. Nishimura, J. Phys. Soc. Japan <u>54</u>, 1224 (1985)
- 19. A. Jain and D. G. Thompson, J. Phys. B 16, 3077 (1983)
- 20. E. Brüche, Ann. Phys. 1, 93, (1929)
- 21. V. F. Sokolv and Y. A. Sokolova, Soviet Tehc. Phys. Letters 7, 268 (1981)
- O. Sueoka, S. Mori and Y. Katayama, J. Phys. B <u>19</u>, L 373, (1986)
- 23. C. Szmytkowski, Chem. Phys. 136, 363 (1987)
- 24. C. Szmytkowski, A. Secca, G. Karwasz and S. Oss, Proceeding of Fifteenth International Conference on the Physics of Electronic and Atomic Collisions, edited by Geddes, Gilbody, Kingston, Latimer and Walters, Brighton, U.K. (1987) P 270
- 25. A. Katase, K. Ishibashi, Y. Matsumoto, T. Sakae, S. Maezono, E. Murakami, K. Watanabe and H. Maki, J. Phys. B 19, 2715 (1986)
- J. L. Pack, R. E. Voshal and A. V. Phelps, Phys. Rev. <u>127</u>, 2084 (1962)
- 27. Y. Itikawa, AT. Data Nucl. Data Tables 21, 69 (1978)
- R. N. Crompton and L. G. Christophorou, Phys. Rev. <u>154</u>, 110, (1967)
- 29. D. S. Belic, M. Landau and R. I. Hall, J. Phys. <u>B 14</u>, 175 (1981)




Figure 1 Total and Partial Ionization Cross Sections of ${\rm H}_2{\rm O}$





Figure 2 Emission Excitation Cross Sections of Dissociative Excitations Products of H $^{\circ}$ and their Sum ($\sigma_{T})$






Figure 4 Excitation Cross Sections of the $\mathrm{H}_2\mathrm{O}$ Vibrational Levels





Figure 5 Rotational Excitation Cross Section of $\rm H_2O$






Figure 7 Elastic Scattering Cross Section in $\rm H_20$ shown along with the Total Scattering Cross Section

DISTRIBUTION LIST

Naval Research Laboratory 4555 Overlook Avenue, S.W. Washington, DC 20375-5000

| | Code | Copies |
|---|--|---|
| Colombant, Dr. Cooperstein, G. Dr. Fernsler, R. Dr. Friedman, M. Dr. Haber, I. Dr. Hubbard, R. F. Dr. Joyce, G. Dr. Lampe, M. Dr. Lau, Y. Y. Dr. Library Meger, M. Dr. Murphy, D. Dr. Ossakow, S. Dr. Pechacek, R. Dr. Picone, M. Dr. Robson, A. Dr. Sprangle, P. Dr. | 4792 4700.1 4791 4792 4792 4790 2628 4750 4751 4700 4750.1 4440 4760 | 30 1 1 1 1 1 1 1 1 2 1 22 1 26 1 1 1 1 |
| Code | 1220 | 1 |

Air Force Office of Scientific Research Physical and Geophysical Sciences Bolling Air Force Base Washington, DC 20332 Attn: Major Bruce Smith

Air Force Weapons Laboratory Kirtland Air Force Base Albuquerque, NM 87117 Attn: W. Baker (AFWL/NTYP)
D. Dietz (AFWL/NTYP) Lt. Col. J. Head

Aberdeen Proving Ground, Maryland 21005
Attn: Dr. Donald Eccleshall (DRXBR-BM)
Dr. Anand Prakash

C. S. Draper Laboratories
555 Technology Square
Cambridge, Massachusetts

Avco Everett Reserch Laboratory 2385 Revere Beach Pkwy Everett, Massachusetts 02149 Attn: Dr. R. Patrick Dr. Dennis Reilly

Ballistic Missile Def. Ad. Tech. Ctr. P.O. Box 1500 Huntsville, Alabama 35807 Attn: Dr. M. Hawie (BMDSATC-1)

Chief of Naval Material Office of Naval Technology MAT-0712, Room 503 800 N. Quincy Street Arlington, VA 22217 Attn: Dr. Eli Zimet

Cornell University 369 Upson Hall Ithaca, NY 14853 Attn: Prof. David Hammer

Department of Energy Washington, DC 20545 Attn: Dr. Wilmot Hess (ET20: GTN, High Energy and Nuclear Physics)

Directed Technologies, Inc. 226 Potomac School Road McLean, VA 22101 Attn: Dr. Ira F. Kuhn Dr. Nancy Chesser

Attn: Dr. E. Olsson Dr. L. Matson

Institute for Fusion Studies University of Texas At Austin RLM 11.218 Austin, Texas 78712 Attn: Prof. Marshall N. Rosenbluth

Intelcom Rad Tech. P.O. Box 81087 San Diego, California 92138 Attn: Dr. W. Selph

Joint Institute for Laboratory Astrophysics National Bureau of Standards and Univ. of Colorado Boulder, CO 80309 Attn: Dr. Arthur V. Phelphs

Kaman Sciences 1500 Garden of the Gods Rd. Colorado Springs, CO 80933 Attn: Dr. John P. Jackson

DASIAC - DETIR Kaman Tempo 25600 Huntington Ave., Suite 500 Alexandria, VA 22303 Attn: Mr. F. Wimenitz

Defense Advanced Res. Projects Agency 1400 Wilson Blvd. Arlington, VA 22209 Attn: Dr. Shen Shey Dr. H. L. Buchanan

Lawrence Livermore Nat'l Laboratory University of California Livermore, California 94550 Attn: Dr. Richard J. Briggs Dr. Simon S. Yu Dr. Frank Chambers

Dr. James W.-K. Mark, L-477

Dr. William Fawley Dr. William Sharp Dr. Daniel S. Prono Dr. John K. Boyd Dr. Kenneth W. Struve Dr. John Clark

Dr. George J. Caporaso Dr. William E. Martin Dr. Donald Prosnitz

Dr. William Barletta

Lockheed Missiles and Space Co. 3251 Hanover Street Bldg. 205, Dept 92-20 Palo Alto, CA 94304 Attn: Dr. John Siambis

Los Alamos Nat'l Scientific Laboratory P.O. Box 1663 Los Alamos, NM 87545

Attn: Dr. L. Thode Dr. M. A. Mostrom, MS-608 Dr. H. Dogliani, MS-5000 Dr. R. Carlson

Ms. Leah Baker, MS-940 Dr. Carl Ekdahl

Lawrence Berkeley Laboratory University of California Berkeley, CA 94720 Attn: Dr. Edward P. Lee

Mission Research Corp. P.O. Drawer 719 Santa Barbara, CA 93102 Attn: Dr. C. Longmire

National Bureau of Standards Gaithersburg, Maryland 20760 Attn: Dr. Mark Wilson

Naval Surface Warfare Center White Oak Laboratory Silver Spring, Maryland 20903 Attn: Dr. R. Cawley Dr. J. W. Forbes

Dr. B. Hui

Mr. W. M. Hinckley Mr. N. E. Scofield

Dr. E. C. Whitman Dr. M. H. Cha Dr. H. S. Uhm

Dr. R. Fiorito

Dr. K. T. Nguyen Dr. R. Stark

Dr. R. Chen

Office of Naval Research 800 North Quincy Street Arlington, VA 22217 Attn: Dr. C. W. Roberson

Office of Naval Research Department of the Navy Code 01231C Arlington, VA 22217 (2 Copies)

Office of Under Secretary of Defense, Research and Engineering Room 3E1034, The Pentagon Washington, DC 20301 Attn: Mr. John M. Bachkosky

Maxwell Laboratories, Inc. 8888 Balboa Avenue San Diego, CA 92123 Attn: Dr. Ken Whitham

McDonnell Douglas Res. Laboratories Dept. 223, Bldg. 33, Level 45 Box 516 St. Louis, MO 63166 Attn: Dr. Evan Rose Dr. Carl Leader

Mission Research Corporation EM Systems Applications 1720 Randolph Road, S.E. Albuquerque, NM 87106 Attn: Dr. Brendan Godfrey Dr. Thomas Hughes Dr. Lawrence Wright

Dr. A. B. Newberger

Physics International, Inc. 2700 Merced Street San Leandro, CA 94577 Attn: Dr. E. Goldman

Princeton University Plasma Physics Laboratory Princeton, NJ 08540 Attn: Dr. Francis Perkins, Jr.

Pulse Science, Inc. 600 McCormack Street San Leandro, CA 94577 Attn: Dr. Sidney Putnam Dr. John Bayless

Sandia National Laboratory Albuquerque, NM 87115 Attn: Dr. Collins Clark Dr. Barbara Epstein

Dr. John Freeman

Dr. Charles Frost Dr. Gordon T. Leifeste

Dr. Gerald N. Hays Dr. James Chang

Dr. Michael G. Mazerakis

Dr. John Wagner Dr. Ron Lipinski

ORI, Inc. 1375 Piccard Drive Rockville, MD 20850 Attn: Dr. C. M. Huddleston

Physical Dynamics, Inc. P.O. Box 1883 La Jolla, California 92038 Attn: Dr. K. Brueckner

Commander Space & Naval Warfare Systems Command, PMW-145 Washington, DC 20363-5100 Attn: CAPT J.D. Fontana CDR W. Bassett

SRI International PSO-15, Molecular Physics Lab 333 Ravenswood Avenue Menlo Park, CA 94025 Attn: Dr. Donald Eckstrom

Strategic Defense Initiative Organization 1717 H Street, N.W. Washington, DC 20009 Attn: Lt. Col. R. L. Gullickson

Titan Systems, Inc. 9191 Towne Centre Dr. -Suite 500 San Diego, CA 92122 Attn: Dr. R. M. Dowe

University of California Physics Department College Park, MD 20742 Attn: Dr. Y. C. Lee Dr. C. Grebogi

University of Michigan Dept. of Nuclear Engineering Ann Arbor, MI 48109 Attn: Prof. Terry Kammash Prof R. Gilgenbach

Science Applications Intl. Corp. P.O. Box 2351 La Jolla, CA 92038 Attn: Dr. Rang Tsang

Science Applications Intl. Corp. 1710 Goodridge Drive McLean, VA 22102 Attn: Mr. W. Chadsey Dr. A. Drobot Dr. K. Papadopoulos Science Applications Intl. Corp 5150 El Camino Road Los Altos, CA 94022 Attn: Dr. R. R. Johnston Dr. Leon Fienstein Dr. Douglas Keeley

Director of Research U.S. Naval Academy Annapolis, MD 21402 (2 Copies)